Nordic Laboratories

PATIENT: XXXXXXXXXXXXXXXXX			TEST REF: TST-NL-XXXX
TEST NUMBER: T-NL-XXXXX (XXXXXXXXXX)		COLLECTED: XX/XX/XXXX	PRACTITIONER: XXXXXXXXXXXXX
GENDER: XYZ		RECEIVED: XX/XX/XXXX	
AGE: XX		TESTED: XX/XX/XXXX	XXXXXXXXXXXXXXXXXXXX

TEST NAME: Metabolomix+ with Urine Elements add-on

3200 Metabolomix+ - FMV Urine

Results Overview



Functional Imbalance Scores

Key: 0-4 : Minimal Need for Support; 5-7 : Moderate Need for Support; 8-10 : High Need for Support

Need for Antioxidant Support	Need for Mitochondrial Support	Need for Inflammation Support	Need for Reduced Exposure	Need for Methylation Support
Oxidative Stress	**Mitochondrial Dysfunction**	**Omega Imbalance**	**Toxic Exposure**	**Methylation Imbalance**
5	0	0	0	0
Cystine ●	Magnesium ▽	Omega-3 Index ●	Lead ●	Methylmalonic Acid ●
Cysteine ▽	FIGLU ●	Omega 6/3 Ratio ▼	Mercury ●	Methionine ▽
Lipid Peroxides △	Methylmalonic Acid ●	α-Linolenic Acid ●	α-Hydroxyisobutyric Acid ●	FIGLU ●
8-OHdG ●	Glutaric Acid △	Arachidonic Acid ▽	α-Ketophenylacetic Acid △	Sarcosine ●
Taurine ▽	Lactic Acid ▽	Linoleic Acid ●	Arsenic ●	Vanilmandelic Acid ●
Citric Acid ▽	Pyruvic Acid ▽	γ-Linolenic Acid ●	Cadmium ●	Arginine ●
cis-Aconitic Acid ▽	Citric Acid ▽	Dihomo-γ-linolenic Acid ●	Pyroglutamic Acid ●	Glycine ●
	cis-Aconitic Acid ▽		Orotic Acid ▽	Serine ●
	Isocitric Acid ●	**OMEGA IMBALANCE NOT INCLUDED IN THIS TEST**	Citric Acid ▽	Creatinine ●
	α-Ketoglutaric Acid ●		cis-Aconitic Acid ▽	
	Succinic Acid ▽		Isocitric Acid ●	
	Malic Acid △		Glutaric Acid △	
	Adipic Acid ●			
	Suberic Acid ●			
	Manganese ●			

Nordic Laboratories Aps
Nygade 6, 3.sal • 1164 Copenhagen K • Denmark
Tlf. +45 33 75 10 00

UK Office:
11 Old Factory Buildings • Stonegate • E. Sussex TN5 7DU • UK
Tel: +44 (0)1580 201 687

www.nordic-labs.com
info@nordic-labs.com

© Copyright 2019 Nordic Laboratories. Reproduction may be made for personal use only. Systematic electronic or print reproduction and distribution including duplication of any material in this paper for a fee or for commercial purposes, or modification of the content of the paper are prohibited.

TEST NAME: Metabolomix+ with Urine Elements add-on

Nutrient Need Overview

	Nutrient Need (0 1 2 3 4 5 6 7 8 9 10)	DRI	Suggested Recommendations	Provider Recommendations
Antioxidants				
Vitamin A		2,333 IU	3,000 IU	
Vitamin C		75 mg	250 mg	
Vitamin E / Tocopherols		22 IU	100 IU	
α-Lipoic Acid			200 mg	
CoQ10			30 mg	
Glutathione				
Plant-based Antioxidants				
B-Vitamins				
Thiamin - B1		1.1 mg	25 mg	
Riboflavin - B2		1.1 mg	25 mg	
Niacin - B3		14 mg	20 mg	
Pyridoxine - B6		1.3 mg	25 mg	
Biotin - B7		30 mcg	200 mcg	
Folate - B9		400 mcg	400 mcg	
Cobalamin - B12		2.4 mcg	500 mcg	
Minerals				
Magnesium		320 mg	600 mg	
Manganese		1.8 mg	3.0 mg	
Molybdenum		45 mcg	75 mcg	
Zinc		8 mg	10 mg	
Essential Fatty Acids				
Omega-3 Fatty Acids		500 mg	500 mg	
GI Support				
Digestive Support/Enzymes			0 IU	
Microbiome Support/Probiotics			10 billion CFU	

Amino Acids (mg/day)

Arginine	0	Methionine	404
Asparagine	187	Phenylalanine	0
Cysteine	108	Serine	0
Glutamine	89	Taurine	837
Glycine	1,277	Threonine	0
Histidine	671	Tryptophan	0
Isoleucine	0	Tyrosine	35
Leucine	0	Valine	0
Lysine	494		

Recommendations for age and gender-specific supplementation are set by comparing levels of nutrient functional need to optimal levels as described in the peer-reviewed literature. They are provided as guidance for short-term support of nutritional deficiencies only.

The Nutrient Need Overview is provided at the request of the ordering practitioner. Any application of it as a therapeutic intervention is to be determined by the ordering practitioner.

© Copyright 2019 Nordic Laboratories. Reproduction may be made for personal use only. Systematic electronic or print reproduction and distribution including duplication of any material in this paper for a fee or for commercial purposes, or modification of the content of the paper are prohibited.

TEST NAME: Metabolomix+ with Urine Elements add-on

Interpretation At-A-Glance

Antioxidant Needs

Vitamin A

- Beta-carotene & other carotenoids are converted to vitamin A (retinol), involved in vision, antioxidant & immune function, gene expression & cell growth.
- Vitamin A deficiency may occur with chronic alcoholism, zinc deficiency, hypothyroidism, or oral contraceptives containing estrogen & progestin.
- Deficiency may result in night blindness, impaired immunity, healing & tissue regeneration, increased risk of infection, leukoplakia or keratosis.
- Food sources include cod liver oil, fortified cereals & milk, eggs, sweet potato, pumpkin, carrot, cantaloupe, mango, spinach, broccoli, kale & butternut squash.

Vitamin C

- Vitamin C is an antioxidant (also used in the regeneration of other antioxidants). It is involved in cholesterol metabolism, the production & function of WBCs and antibodies, and the synthesis of collagen, norepinephrine and carnitine.
- Deficiency may occur with oral contraceptives, aspirin, diuretics or NSAIDs.
- Deficiency can result in scurvy, swollen gingiva, periodontal destruction, loose teeth, sore mouth, soft tissue ulcerations, or increased risk of infection.
- Food sources include oranges, grapefruit, strawberries, tomato, sweet red pepper, broccoli and potato.

Vitamin E / Tocopherols



- Alpha-tocopherol (body's main form of vitamin E) functions as an antioxidant, regulates cell signaling, influences immune function and inhibits coagulation.
- Deficiency may occur with malabsorption, cholestyramine, colestipol, isoniazid, orlistat, olestra and certain anti-convulsants (e.g., phenobarbital, phenytoin).
- Deficiency may result in peripheral neuropathy, ataxia, muscle weakness, retinopathy, and increased risk of CVD, prostate cancer and cataracts.
- Food sources include oils (olive, soy, corn, canola, safflower, sunflower), eggs, nuts, seeds, spinach, carrots, avocado, dark leafy greens and wheat germ.

α-Lipoic Acid

- α-Lipoic acid plays an important role in energy production, antioxidant activity (including the regeneration of vitamin C and glutathione), insulin signaling, cell signaling and the catabolism of α-keto acids and amino acids.
- High biotin intake can compete with lipoic acid for cell membrane entry.
- Optimal levels of α-lipoic acid may improve glucose utilization and protect against diabetic neuropathy, vascular disease and age-related cognitive decline.
- Main food sources include organ meats, spinach and broccoli. Lesser sources include tomato, peas, Brussels sprouts and brewer's yeast.

CoQ10

- CoQ10 is a powerful antioxidant that is synthesized in the body and contained in cell membranes. CoQ10 is also essential for energy production & pH regulation.
- CoQ10 deficiency may occur with HMG-CoA reductase inhibitors (statins), several anti-diabetic medication classes (biguanides, sulfonylureas) or beta-blockers.
- Low levels may aggravate oxidative stress, diabetes, cancer, congestive heart failure, cardiac arrhythmias, gingivitis and neurologic diseases.
- Main food sources include meat, poultry, fish, soybean, canola oil, nuts and whole grains. Moderate sources include fruits, vegetables, eggs and dairy.

Glutathione

- Glutathione (GSH) is composed of cysteine, glutamine & glycine. GSH is a source of sulfate and plays a key role in antioxidant activity and detoxification of toxins.
- GSH requirement is increased with high-fat diets, cigarette smoke, cystinuria, chronic alcoholism, chronic acetaminophen use, infection, inflammation and toxic exposure.
- Deficiency may result in oxidative stress & damage, impaired detoxification, altered immunity, macular degeneration and increased risk of chronic illness.
- Food sources of GSH precursors include meats, poultry, fish, soy, corn, nuts, seeds, wheat germ, milk and cheese.

Plant-based Antioxidants



- Oxidative stress is the imbalance between the production of free radicals and the body's ability to readily detoxify these reactive species and/or repair the resulting damage with anti-oxidants.
- Oxidative stress can be endogenous (energy production and inflammation) or exogenous (exercise, exposure to environmental toxins).
- Oxidative stress has been implicated clinically in the development of neurodegenerative diseases, cardiovascular diseases and chronic fatigue syndrome.
- Antioxidants may be found in whole food sources (e.g., brightly colored fruits & vegetables, green tea, turmeric) as well as nutraceuticals (e.g., resveratrol, EGCG, lutein, lycopene, ginkgo, milk thistle, etc.).

KEY

Function of Nutrient | Cause of Deficiency | Complications of Deficiency | Food Sources of Nutrient

© Copyright 2019 Nordic Laboratories. Reproduction may be made for personal use only. Systematic electronic or print reproduction and distribution including duplication of any material in this paper for a fee or for commercial purposes, or modification of the content of the paper are prohibited.

TEST NAME: Metabolomix+ with Urine Elements add-on

Interpretation At-A-Glance

B-Vitamin Needs

Thiamin - B1

- B1 is a required cofactor for enzymes involved in energy production from food, and for the synthesis of ATP, GTP, DNA, RNA and NADPH.
- Low B1 can result from chronic alcoholism, diuretics, digoxin, oral contraceptives and HRT, or large amounts of tea & coffee (contain anti-B1 factors).
- B1 deficiency may lead to dry beriberi (e.g., neuropathy, muscle weakness), wet beriberi (e.g., cardiac problems, edema), encephalopathy or dementia.
- Food sources include lentils, whole grains, wheat germ, Brazil nuts, peas, organ meats, brewer's yeast, blackstrap molasses, spinach, milk & eggs.

Riboflavin - B2

- B2 is a key component of enzymes involved in antioxidant function, energy production, detoxification, methionine metabolism and vitamin activation.
- Low B2 may result from chronic alcoholism, some anti-psychotic medications, oral contraceptives, tricyclic antidepressants, quinacrine or adriamycin.
- B2 deficiency may result in oxidative stress, mitochondrial dysfunction, low uric acid, low B3 or B6, high homocysteine, anemia or oral & throat inflammation.
- Food sources include milk, cheese, eggs, whole grains, beef, chicken, wheat germ, fish, broccoli, asparagus, spinach, mushrooms and almonds.

Niacin - B3

- B3 is used to form NAD and NADP, involved in energy production from food, fatty acid & cholesterol synthesis, cell signaling, DNA repair & cell differentiation.
- Low B3 may result from deficiencies of tryptophan (B3 precursor), B6, B2 or Fe (cofactors in B3 production), or from long-term isoniazid or oral contraceptive use.
- B3 deficiency may result in pellagra (dermatitis, diarrhea, dementia), neurologic symptoms (e.g., depression, memory loss), bright red tongue or fatigue.
- Food sources include poultry, beef, organ meats, fish, whole grains, peanuts, seeds, lentils, brewer's yeast and lima beans.

Pyridoxine - B6

- B6 (as P5P) is a cofactor for enzymes involved in glycogenolysis & gluconeogenesis, and synthesis of neurotransmitters, heme, B3, RBCs and nucleic acids.
- Low B6 may result from chronic alcoholism, long-term diuretics, estrogens (oral contraceptives and HRT), anti-TB meds, penicillamine, L-DOPA or digoxin.
- B6 deficiency may result in neurologic symptoms (e.g., irritability, depression, seizures), oral inflammation, impaired immunity or increased homocysteine.
- Food sources include poultry, beef, beef liver, fish, whole grains, wheat germ, soybean, lentils, nuts & seeds, potato, spinach and carrots.

Biotin - B7

- Biotin is a cofactor for enzymes involved in functions such as fatty acid synthesis, mitochondrial FA oxidation, gluconeogenesis and DNA replication & transcription.
- Deficiency may result from certain inborn errors, chronic intake of raw egg whites, long-term TPN, anticonvulsants, high-dose B5, sulfa drugs & other antibiotics.
- Low levels may result in neurologic symptoms (e.g., paresthesias, depression), hair loss, scaly rash on face or genitals or impaired immunity.
- Food sources include yeast, whole grains, wheat germ, eggs, cheese, liver, meats, fish, wheat, nuts & seeds, avocado, raspberries, sweet potato and cauliflower.

Folate - B9

- Folate plays a key role in coenzymes involved in DNA and SAMe synthesis, methylation, nucleic acids & amino acid metabolism and RBC production.
- Low folate may result from alcoholism, high-dose NSAIDs, diabetic meds, H2 blockers, some diuretics and anti-convulsants, SSRIs, methotrexate, trimethoprim, pyrimethamine, triamterene, sulfasalazine or cholestyramine.
- Folate deficiency can result in anemia, fatigue, low methionine, increased homocysteine, impaired immunity, heart disease, birth defects and CA risk.
- Food sources include fortified grains, green vegetables, beans & legumes.

Cobalamin - B12

- B12 plays important roles in energy production from fats & proteins, methylation, synthesis of hemoglobin & RBCs, and maintenance of nerve cells, DNA & RNA.
- Low B12 may result from alcoholism, malabsorption, hypochlorhydria (e.g., from atrophic gastritis, H. pylori infection, pernicious anemia, H2 blockers, PPIs), vegan diets, diabetic meds, cholestyramine, chloramphenicol, neomycin or colchicine.
- B12 deficiency can lead to anemia, fatigue, neurologic symptoms (e.g., paresthesias, memory loss, depression, dementia), methylation defects or chromosome breaks.
- Food sources include shellfish, red meat, poultry, fish, eggs, milk and cheese.

KEY

Function of Nutrient | Cause of Deficiency | Complications of Deficiency | Food Sources of Nutrient

© Copyright 2019 Nordic Laboratories. Reproduction may be made for personal use only. Systematic electronic or print reproduction and distribution including duplication of any material in this paper for a fee or for commercial purposes, or modification of the content of the paper are prohibited.

TEST NAME: Metabolomix+

Interpretation At-A-Glance

Mineral Needs

Magnesium

- Magnesium is involved in >300 metabolic reactions. Key areas include energy production, bone & ATP formation, muscle & nerve conduction and cell signaling.
- Deficiency may occur with malabsorption, alcoholism, hyperparathyroidism, renal disorders (wasting), diabetes, diuretics, digoxin or high doses of zinc.
- Low Mg may result in muscle weakness/spasm, constipation, depression, hypertension, arrhythmias, hypocalcemia, hypokalemia or personality changes.
- Food sources include dark leafy greens, oatmeal, buckwheat, unpolished grains, chocolate, milk, nuts & seeds, lima beans and molasses.

Manganese

- Manganese plays an important role in antioxidant function, gluconeogenesis, the urea cycle, cartilage & bone formation, energy production and digestion.
- Impaired absorption of Mn may occur with excess intake of Fe, Ca, Cu, folic acid, or phosphorous compounds, or use of long-term TPN, Mg-containing antacids or laxatives.
- Deficiency may result in impaired bone/connective tissue growth, glucose & lipid dysregulation, infertility, oxidative stress, inflammation or hyperammonemia.
- Food sources include whole grains, legumes, dried fruits, nuts, dark green leafy vegetables, liver, kidney and tea.

Molybdenum

- Molybdenum is a cofactor for enzymes that convert sulfites to sulfate, and nucleotides to uric acid, and that help metabolize aldehydes & other toxins.
- Low Mo levels may result from long-term TPN that does not include Mo.
- Mo deficiency may result in increased sulfite, decreased plasma uric acid (and antioxidant function), deficient sulfate, impaired sulfation (detoxification), neurologic disorders or brain damage (if severe deficiency).
- Food sources include buckwheat, beans, grains, nuts, beans, lentils, meats and vegetables (although Mo content of plants depends on soil content).

Zinc

- Zinc plays a vital role in immunity, protein metabolism, heme synthesis, growth & development, reproduction, digestion and antioxidant function.
- Low levels may occur with malabsorption, alcoholism, chronic diarrhea, diabetes, excess Cu or Fe, diuretics, ACE inhibitors, H2 blockers or digoxin.
- Deficiency can result in hair loss and skin rashes, also impairments in growth & healing, immunity, sexual function, taste & smell and digestion.
- Food sources include oysters, organ meats, soybean, wheat germ, seeds, nuts, red meat, chicken, herring, milk, yeast, leafy and root vegetables.

KEY

Function of Nutrient | Cause of Deficiency | Complications of Deficiency | Food Sources of Nutrient

© Copyright 2019 Nordic Laboratories. Reproduction may be made for personal use only. Systematic electronic or print reproduction and distribution including duplication of any material in this paper for a fee or for commercial purposes, or modification of the content of the paper are prohibited.

TEST NAME: Metabolomix+ with Urine Elements add-on

Interpretation At-A-Glance

Microbiome & Digestive Support

Microbiome Support/Probiotics



- Probiotics have many functions. These include: production of some B vitamins and vitamin K; enhance digestion & absorption; decrease severity of diarrheal illness; modulate of immune function & intestinal permeability.
- Alterations of gastrointestinal microflora may result from C-section delivery, antibiotic use, improved sanitation, decreased consumption of fermented foods and use of certain drugs.
- Some of the diseases associated with microflora imbalances include: IBS, IBD, fibromyalgia, chronic fatigue syndrome, obesity, atopic illness, colic and cancer.
- Food sources rich in probiotics are yogurt, kefir and fermented foods.

Digestive Support/Enzymes



- Pancreatic enzymes are secreted by the exocrine glands of the pancreas and include protease/peptidase, lipase and amylase.
- Pancreatic exocrine insufficiency may be primary or secondary in nature. Any indication of insufficiency warrants further evaluation for underlying cause (i.e., celiac disease, small intestine villous atrophy, small bowel bacterial overgrowth).
- A high functional need for digestive enzymes suggests that there is an impairment related to digestive capacity.
- Determining the strength of the pancreatic enzyme support depends on the degree of functional impairment. Supplement potency is based on the lipase units present in both prescriptive and non-prescriptive agents.

Functional Imbalances

Mitochondrial Dysfunction



- Mitochondria are a primary site of generation of reactive oxygen species. Oxidative damage is considered an important factor in decline of physiologic function that occurs with aging and stress.
- Mitochondrial defects have been identified in cardiovascular disease, fatigue syndromes, neurologic disorders such as Parkinson's and Alzheimer's disease, as well as a variety of genetic conditions. Common nutritional deficiencies can impair mitochondrial efficiency.

Need for Methylation

- Methylation is an enzymatic process that is critical for both synthesis and inactivation. DNA, estrogen and neurotransmitter metabolism are all dependent on appropriate methylation activity.
- B vitamins and other nutrients (methionine, magnesium, selenium) functionally support catechol-O-methyltransferase (COMT), the enzyme responsible for methylation.

Toxic Exposure

- Methyl tert-Butyl Ether (MTBE) is a common gasoline additive used to increase octane ratings, and has been found to contaminate ground water supplies where gasoline is stored. Inhalation of MTBE may cause nose and throat irritation, as well as headaches, nausea, dizziness and mental confusion. Animal studies suggest that drinking MTBE may cause gastrointestinal irritation, liver and kidney damage and nervous system effects.
- Styrene is classified by the US EPA as a "potential human carcinogen," and is found widely distributed in commercial products such as rubber, plastic, insulation, fiberglass, pipes, food containers and carpet backing.
- Levels of these toxic substances should be examined within the context of the body's functional capacity for methylation and need for glutathione.

KEY

Function of Nutrient | Cause of Deficiency | Complications of Deficiency | Food Sources of Nutrient

© Copyright 2019 Nordic Laboratories. Reproduction may be made for personal use only. Systematic electronic or print reproduction and distribution including duplication of any material in this paper for a fee or for commercial purposes, or modification of the content of the paper are prohibited.

TEST NAME: Metabolomix+ with Urine Elements add-on

Oxidative Stress & Mitochondrial Dysfunction



© Copyright 2019 Nordic Laboratories. Reproduction may be made for personal use only. Systematic electronic or print reproduction and distribution including duplication of any material in this paper for a fee or for commercial purposes, or modification of the content of the paper are prohibited.

TEST NAME: Metabolomix+ with Urine Elements add-on

All biomarkers reported in mmol/mol creatinine unless otherwise noted.

Organic Acids

Malabsorption & Dysbiosis Markers

Malabsorption Markers

		Reference Range
Indoleacetic Acid	0.6	<= 4.2
Phenylacetic Acid	0.04	<= 0.12

Dysbiosis Markers

		Reference Range
Dihydroxyphenylpropionic Acid (DHPPA)	0.3	<= 5.3
3-Hydroxyphenylacetic Acid	0.4	<= 8.1
4-Hydroxyphenylacetic Acid	2	<= 29
Benzoic Acid	0.05	<= 0.05
Hippuric Acid	1	<= 603

Yeast / Fungal Dysbiosis Markers

		Reference Range
D-Arabinitol	16	<= 36
Citramalic Acid	0.4	<= 5.8
Tartaric Acid	1	<= 15

Cellular Energy & Mitochondrial Markers

Fatty Acid Metabolism

		Reference Range
Adipic Acid	1.9	<= 2.8
Suberic Acid	0.3	<= 2.1

Carbohydrate Metabolism

		Reference Range
Pyruvic Acid	10	7-32
Lactic Acid	3.2	1.9-19.8
α-Hydroxybutyric Acid	0.60	<= 0.83
β-OH-Butyric Acid	0.5	<= 2.8
β-OH-β-Methylglutaric Acid	1	<= 15

Energy Metabolism

		Reference Range
Citric Acid	94	40-520
cis-Aconitic Acid	11	10-36
Isocitric Acid	44	22-65
α-Ketoglutaric Acid	10	4-52
Succinic Acid	0.5	0.4-4.6
Malic Acid	2.1	<= 3.0

Methodology: GCMS, LC/MS/MS, Alkaline Picrate, Colorimetric

Vitamin Markers

Branched-Chain Catabolites (B1, B2, B3, ALA)

		Reference Range
α-Ketoadipic Acid	0.4	<= 1.7
α-Ketoisovaleric Acid	0.24	<= 0.97
α-Ketoisocaproic Acid	0.87	<= 0.89
α-Keto-β-Methylvaleric Acid	0.4	<= 2.1
Glutaric Acid	0.43	<= 0.51
Isovalerylglycine	0.4	<= 3.7

Methylation Markers (Folate, B12)

		Reference Range
Formiminoglutamic Acid (FIGlu)	0.7	<= 1.5
Methylmalonic Acid	0.5	<= 1.9

Biotin Markers

		Reference Range
3-Hydroxypropionic Acid	16	5-22
3-Hydroxyisovaleric Acid	2	<= 29

Neurotransmitter Metabolites

Kynurenine Markers (Vitamin B6)

		Reference Range
Kynurenic Acid	0.3	<= 7.1
Quinolinic Acid	0.3	<= 9.1
Kynurenic / Quinolinic Ratio	1.00	>= 0.44
Xanthurenic Acid	0.28	<= 0.96

Catecholamine Markers

		Reference Range
Homovanillic Acid	1.6	1.2-5.3
Vanilmandelic Acid	1.4	0.4-3.6
3-Methyl-4-OH-phenylglycol	0.15	0.02-0.22

Serotonin Markers

		Reference Range
5-OH-indoleacetic Acid	4.5	3.8-12.1

Toxin & Detoxification Markers

		Reference Range
Pyroglutamic Acid	26	16-34
α-Ketophenylacetic Acid (from Styrene)	0.38	<= 0.46
α-Hydroxyisobutyric Acid (from MTBE)	0.5	<= 6.7
Orotic Acid	0.36	0.33-1.01

Organic Acid Reference Ranges are Age Specific

© Copyright 2019 Nordic Laboratories. Reproduction may be made for personal use only. Systematic electronic or print reproduction and distribution including duplication of any material in this paper for a fee or for commercial purposes, or modification of the content of the paper are prohibited.

TEST NAME: Metabolomix+ with Urine Elements add-on

Methodology: Colorimetric, thiobarbituric acid reactive substances (TBARS), Alkaline Picrate, Hexokinase/G-6-PDH, HPLC, GC/MS

Organic Acids

Oxalate Markers		Reference Range
Glyceric Acid	12.2	3.5-16.4
Glycolic Acid	14	<= 67
Oxalic Acid	65	<= 78

All biomarkers reported in mmol/mol creatinine.

Creatinine Concentration		Reference Range
Creatinine ◆	8.8	3.1-19.5 mmol/L

Oxidative Stress Markers

Oxidative Damage		Reference Range
Lipid Peroxides (urine)	8.3	<= 10.0 micromol/g Creat.
8-OHdG (urine)	5	<= 15 mcg/g Creat.

The Oxidative Stress reference ranges are based on an adult population.

Pathways

Methylation Markers





Branch-Chain Amino Acid Metabolism



© Copyright 2019 Nordic Laboratories. Reproduction may be made for personal use only. Systematic electronic or print reproduction and distribution including duplication of any material in this paper for a fee or for commercial purposes, or modification of the content of the paper are prohibited.

TEST NAME: Metabolomix+ with Urine Elements add-on

All biomarkers reported in micromol/g creatinine unless otherwise noted.

Amino Acids (FMV)

Nutritionally Essential Amino Acids

Amino Acid		Reference Range
Arginine	19	3-43
Histidine	163	124-894
Isoleucine	19	3-28
Leucine	26	4-46
Lysine	29	11-175
Methionine	3	2-18
Phenylalanine	23	8-71
Taurine	31	21-424
Threonine	69	17-135
Tryptophan	19	5-53
Valine	33	7-49

Nonessential Protein Amino Acids

Amino Acid		Reference Range
Alanine	63	63-356
Asparagine	40	25-166
Aspartic Acid	13	<= 14
Cysteine	16	8-74
Cystine	19	10-104
γ-Aminobutyric Acid	3	<= 5
Glutamic Acid	15	4-27
Glutamine	188	110-632
Proline	6	1-13
Tyrosine	30	11-135

Creatinine Concentration

		Reference Range
Creatinine◆	7.1	3.1-19.5 mmol/L

Amino Acid reference ranges are age specific.

Methodology: LC/MS/MS, Alkaline Picrate

Intermediary Metabolites

B-Vitamin Markers		Reference Range
α-Aminoadipic Acid	19	2-47
α-Amino-N-butyric Acid	15	2-25
β-Aminoisobutyric Acid	16	11-160
Cystathionine	15	2-68

Urea Cycle Markers		
Citrulline	1.3	0.6-3.9
Ornithine	15	2-21
Urea◆	357	168-465 mmol/g creatinine

Glycine/Serine Metabolites		
Glycine	138	95-683
Serine	69	40-163
Ethanolamine	73	50-235
Phosphoethanolamine	4	1-13
Phosphoserine	6	3-13
Sarcosine	0.5	<= 1.1

Dietary Peptide Related Markers		Reference Range
Anserine (dipeptide)	18.8	0.4-105.1
Carnosine (dipeptide)	15	1-28
1-Methylhistidine	45	38-988
3-Methylhistidine	50	44-281
β-Alanine	15	<= 22

© Copyright 2019 Nordic Laboratories. Reproduction may be made for personal use only. Systematic electronic or print reproduction and distribution including duplication of any material in this paper for a fee or for commercial purposes, or modification of the content of the paper are prohibited.

TEST NAME: Metabolomix+ with Urine Elements add-on

3204 Add - on Comprehensive Urine Elements - FMV Urine

Methodology: ICP-MS and Alkaline Picrate

Elemental Markers

Toxic Elements

Element	Results in ug/g creatinine	Reference Range
Lead	1.0	<= 1.4
Mercury	0.50	<= 2.19
Aluminum	2.5	<= 22.3
Antimony	0.025	<= 0.149
Arsenic	1	<= 50
Barium	0.5	<= 6.7
Bismuth	0.75	<= 2.28
Cadmium	0.25	<= 0.64
Cesium	2.5	<= 10.5
Gadolinium	0.010	<= 0.019
Gallium	0.003	<= 0.028
Nickel	2.50	<= 3.88
Niobium	0.040	<= 0.084
Platinum	0.015	<= 0.033
Rubidium	3	<= 2,263
Thallium	0.050	<= 0.298
Thorium	2.500	<= 4.189
Tin	0.10	<= 2.04
Tungsten	0.025	<= 0.211
Uranium	0.010	<= 0.026

Nutrient Elements

Element	Results in ug/g creatinine	Reference Range
Chromium	2.5	0.6-9.4
Cobalt	2.50	0.01-2.60
Copper	5.0	4.0-11.4
Iron	10	5-64
Lithium	25	9-129
Manganese	0.25	0.03-1.16
Molybdenum	25	15-175
Selenium	180	32-333
Strontium	125	47-346
Vanadium	2.5	0.1-3.2
Zinc	250	63-688

Element	Results in mg/g creatinine	Reference Range
Calcium	40	37-313
Magnesium	48	41-267
Potassium	2,550	759-4,653
Sulfur	390	367-1,328

Creatinine Concentration

		Reference Range
Creatinine ◆	100.00	23.00-205.00 mmol/L

The performance characteristics of all assays have been verified by Genova Diagnostics, Inc. Unless otherwise noted with ◆, the assays have not been cleared by the U.S. Food and Drug Administration.

© Copyright 2019 Nordic Laboratories. Reproduction may be made for personal use only. Systematic electronic or print reproduction and distribution including duplication of any material in this paper for a fee or for commercial purposes, or modification of the content of the paper are prohibited.